Public Meeting Information

Asian Citrus Psyllid and Huanglongbing Response Program



Asian Citrus Psyllid (ACP)
Life Cycle

- Up to 10 generations per year
- 5 nymph stages
- Life cycle is 15 47 days
- Eggs can reach the adult stage in
 2 weeks

Life Stages



Eggs



Nymphs with waxy tubules




Adult

Adult feeding with nymphs



Huanglongbing (HLB)







Asymmetric leaf mottling



Blotchy mottling



Yellow, corked veins

- HLB is a bacterial disease of citrus and citrus relatives, caused by the bacteria Candidatus Liberibacter asiaticus.
- HLB can be spread by ACP feeding or grafting with infected budwood. ACP ingest the bacteria when feeding and can spread it to other trees.
- Symptoms include yellow shoots, asymmetric leaf mottle, thick or woody leaf veins, and lopsided, inedible fruit.
- There is no known cure for HLB. Once a tree is infected with HLB, it will die.
- If you suspect you have a citrus tree with HLB, please contact the California Department of Food and Agriculture at 1-800-491-1899.



Survey


CDFA staff will survey the area for all host plants and look for ACP and HLB disease symptoms.








- CDFA surveyors will collect leaf samples and any ACP life stages from all host plants.
- Host plants include all citrus varieties and hybrids (kumquat, mandarin, grapefruit, orange, lemon, and lime) and closely related plants like curry leaf and ornamental orange-jasmine.
- All plant samples are shipped to the CDFA Plant Pest Diagnostics Center laboratory in Sacramento and ACP samples are shipped to the Citrus Research Board laboratory in Riverside for analysis.
- If a sample is confirmed positive for ACP or the HLB disease, the resident will be notified.





Treatment

- CDFA will be treating citrus trees to eradicate possibly infected ACP in the area.
- Treatment will be applied by a professional applicator or CDFA staff.









Active Ingredients

- The ACP treatment program uses products containing the active ingredients imidacloprid and beta-cyfluthrin.
- Imidacloprid is an insecticide that provides lasting protection to trees. It is available in a variety of home garden products and flea treatments for dogs and cats.
- Beta-cyfluthrin is an insecticide used for the control of ACP adults and nymphs. It is similar to a natural compound found in chrysanthemum flowers. Beta-cyfluthrin products are used in homes, restaurants, hospitals, food processing plants and gardens.



Tempo SC Ultra Beta-cyfluthrin product




Merit 2F Imidacloprid product




CoreTect Imidacloprid product



Notifications



48-hour Pre-treatment notification



Reschedule notification

	CALIFORNIA DEPARTM COUNTY AGRICULTU	ENT OF FOOD AND AG IRAL COMMISSIONER	
	NOTICE OF INS	SECTICIDE TREATM	MENT
	Department of Food and Agricu pest, or Huanglongbing (HLB), a		the Asian citrus psyllid (ACP), a your neighborhood.
citrus trees or which poses	your property, you have made a	low. By allowing CDFA to a huge contribution to the s 's environment and econo	A.M. / P.M., your property perform emergency treatment of suppression of this invasive pest, omy. If you have any questions,
		the ACP and HLB, please o://www.cdfa.ca.gov/phpp	
	insecticide for controlling Asia	in citrus psyllid adults an	ion of a pyrethroid contact d nymphs. This material will be plants on designated residential
	designated residential propertie	es. The material is taken	to the soil of all host plants at up into the plant via the root ion against the pest, depending
	soil surface of host plants at d	esignated residential prop em and provides approxi	blets were applied below the lecties. The material is taken up mately 12 months of protection
CDFA has sel environmenta	ected the insecticide products at safety, and California registratio	ove based on their effection status.	veness against ACP, worker and
•	Once the area has dried, you m	ay use your property as yo	u normally would.
•	To prevent the spread of this pe property without contacting our	st, please do not move pla project office.	nt material off your
	For best results, please irrigate	the plants that were treater	d within 24 hours of treatment.
	are experiencing health problems at (800) 222-1222(voice), or co		the California Poison Control
Thank you fo	or your cooperation.		
BLOCK NUN (Please mention the bit 80-234 (3/14/18)	BER_ book number when contacting our office)	DATE	
Pest Detection/Emerge	ncy Projects 999 • www.cdfa.ca.gov		State of California Gavin Newsom, Governor

Post-treatment notification



Pest Hotline: 1.800.491.1899

CITRUS PEST & DISEASE PREVENTION DIVISION CALIFORNIA DEPARTMENT OF FOOD AND AGRICULTURE

Tree Removal

- There is no cure for HLB; once a tree is infected it will die.
- If an HLB positive tree is confirmed, the tree will be removed to prevent spread of the HLB disease to other citrus trees.









Quarantine

- ACP and HLB quarantines help prevent the spread of citrus pests.
- Avoid sharing fruit, if possible.
 - Homegrown fruit without stems and leaves maybe moved within the quarantine (25 pounds or less).
- Do not move potted citrus plants, budwood, cuttings, and other host plants from your property.
- Only graft using plant parts received from an approved source.
- Only nurseries with a CDFA approved structure may sell citrus plants with a red HLB quarantine tag.
- Contact CDFA for questions about moving citrus plants or fruit.





www.cdfa.ca.gov/citrus/pests_diseases /hlb/regulation





Biological Control

- Tamarixia radiata is an effective biocontrol agent and a natural enemy of ACP.
- This method alone is not enough to stop the spread of ACP.
- Strategic releases, along with treatment applications, help suppress ACP.
- Tamarixia are not a threat to humans or pets.



ACP Mummy



Tamarixia radiata



Tamarixia and ACP



Tamarixia laying eggs on an ACP nymph



Protecting Pollinators

- Protecting native bees and managed hives is essential to maintaining our food supply and the environment.
- CDFA staff actively look for foraging bees and take precautions to prevent pesticide drift.
- The program follows all pesticide labels and instructions for bee safety.
- CDFA will not treat if bees are present.








More info online at www.cdfa.ca.gov/plant/pollinators



Environmental Consultation and Monitoring


- Consultations are conducted with other State agencies prior to beginning any activity.
- Monitoring is conducted when activities move into new counties.
 - Air, soil, leaf and fruit samples are collected and tested for pesticide levels before and after applications.







https://californiacitrusthreat.org







A Threat to California Citrus

A plant disease that kills citrus trees has been found in California. The disease, called Huanglongbing or citrus greening disease, isn't harmful to humans, but it is fatal for citrus trees and has no cure. The disease is spread by a pest called the Asian citrus psyllid as it feeds on citrus tree leaves. Until researchers find a solution, California homeowners who enjoy growing fresh citrus fruit in their yards, and




PEST & DISEASE PROTECT YOUR CITRUS CALIFORNIA CITRUS NEWS RESOURCES

Public Meetings

In addition to being available on <u>CDFA's website</u>, presentations for recent public meetings are available on <u>CaliforniaCitrusThreat.org</u>. Presentations include information the pest and disease, upcoming ac agricultural officials in specific communities and regulations in place to limit the spread of the ps

Learn More








http://ipm.ucanr.edu/







https://www.cdfa.ca.gov/citrus/



CDFA Home | Citrus Pest and Disease Prevention Division

Citrus Pest and Disease Prevention Division

1220 "N" Street, Sacramento, CA 95814 • 916-274-6300 • cdfa.cpdpd@cdfa.ca.gov

A Pest Hotline: 1-800-491-1899

The California Department of Food and Agriculture (CDFA) established the Citrus Pest and Disease Prevention Program (CPDPP) in 2009 to sustain and protect California citrus in accordance with the Food and Agriculture Code, section 5911-5940. In response to increasing pest and disease pressure, the Citrus Pest and Disease Prevention Committee recommended that CDFA seek dedicated resources to implement the CPDPP. Dedicated resources were secured in the 2019 Budget Act and the Citrus Pest and Disease Prevention Division (CPDPD) was established in July 2019. CPDPP activities previously carried out by the CDFA Plant Health and Pest Prevention Services Division have been transitioned



Hot Topics

- What to Expect when CDFA is Inspecting, Trapping and Treating
- ▶ Health Questions and Answers CoreTect®
- ▶ Health Questions and Answers Merit® 2F
- ▶ Health Questions and Answers Tempo® SC Ultra
- Analyses regarding the state of ACP and HLB in California





Thank you for attending!

- https://californiacitrusthreat.org
- http://ipm.ucanr.edu/
- https://www.cdfa.ca.gov/citrus/
- https://www.cdfa.ca.gov/plant/reportapest/

